

# Energy Balance of the Climate System: Contribution of Clouds and Aerosol to Uncertain Futures



James C. Wilson  
Mechanical and Materials Engineering  
University of Denver

# Energy Balance of the Climate System: Contribution of Clouds and Aerosol to Uncertain Futures



This document includes some changes made to improve clarity and add some additional information beyond that presented in the talk


James C. Wilson  
Mechanical and Materials Engineering  
University of Denver

# Acknowledgments:

William Anders, Jim Lovell, Frank Borman

Joseph Fourier


Daniel Murphy




Viewing and Reading;

[www.nytimes.com/video/opinion/100000005831656/earthrise.html](http://www.nytimes.com/video/opinion/100000005831656/earthrise.html)

Murphy, et al., "An observationally based energy balance for Earth...", *JGR*, 114, 2009, <https://doi.org/10.1029/2009JD012105>



1. Climate system gets its energy from the sun
2. Climate system gets rid of energy by radiating infrared (IR) to space. Fourier called it dark energy after Hershel's 1800 experiment.
3. Absorbed solar > escaping IR -> Warming
4. Escaping IR > Absorbed solar-> Cooling




Vertical Axis:  
Watts/m<sup>2</sup> – μm


The climate radiates  
in the infrared  
because of its  
temperature.

Wavelength  
differences result in  
different interactions  
and fates in the  
atmosphere.

Fig:  
Andy Dessler



# ImBalance: Trenberth, Fasullo and Kiehl, BAMS, 2009




Imbalance is 0.9 Watts/m<sup>2</sup> and is small compared to incoming and outgoing. It is inferred from ocean from warming.


GHGs impede the escape of upwelling IR and add to downwelling energy, Water vapor is a potent GHG controlled by temperature and increased by warming driven by increases in CO<sub>2</sub>, etc ,

FIG. 1. The global annual mean Earth's energy budget for the Mar 2000 to May 2004 period ( $W m^{-2}$ ). The broad arrows indicate the schematic flow of energy in proportion to their importance.

# GLOBAL ENERGY BALANCE (Unit: $\text{W}\cdot\text{m}^{-2}$ )

1987: Satellite data permitted quantitative description of the energy balance 30 years ago.





**Figure 1.** Illustrative sketch of the response of the Earth energy budget to an idealized step function change in a greenhouse gas. Outgoing radiation from the Earth at first decreases then returns toward an equilibrium value as the Earth warms. An initial rapid temperature response, as plotted here, reduces the amount of energy available to heat the Earth and slows the eventual approach to equilibrium.

Murphy et al, 2009

1. Start in balance

2. Add GHG -

Escaping Infrared  
is reduced


3. Planet warms &  
escaping IR

increases ( $\sigma T^4$ )

5. Balance



restored at warmer  
temperature.

# GAST is going up. But it is not associated with a defined mass. So we can't apply First Law to it.




# Added GHG's Impede Escape of IR and add Energy to the climate system.

Murphy et al, 2009. IPCC



# The Change in Outgoing IR...


- Has been validated some time ago with spectrometers on satellites
  - The amounts of energy and the spectral details.
  - <https://www.nature.com/articles/35066553>




# To Summarize

- Oceans get almost all of added energy.
- Added GHGs have reduced outgoing radiation by ~1700 ZJ (over plotted time period)
- 275 ZJ have showed up in climate
- Warming is the effect. Added GHGs are a cause.
- We have about 5x the cause that we need to explain the effect.
- If another cause is to be proposed, the entire 1700 ZJ from GHGs must be disposed of without warming climate and 275 ZJ must be found elsewhere. Also observed changes in outgoing IR intensity and wavelength need another explanation.

# To Summarize

- So why haven't we warmed more?
  - The answer is in changes in clouds and aerosols.
  - The amounts are shown as a difference in the next slide.
  - The increases in outgoing radiation minus the impact of the water vapor feedback are shown in the pink band. They reduce the amount of cooling required of aerosols and clouds.
- 

Daniel Murphy: NOAA. Updated.



# Aerosol Effects have been growing. So have:

- Sulfur Emissions (Direct Effect)

$\text{SO}_2 \rightarrow \text{H}_2\text{SO}_4 \rightarrow \text{Particles} \rightarrow \text{light reflected to space}$

- Cloud Modification (Indirect Effect)

- Clouds form on small particles



- Small particles have increased in number

- More cloud droplets reflect more light to space


- Other effects as well.

# Model Projections of Future Temperatures: Range of projected temperatures shown by the width of the colored for each scenario.

Greater Emissions Lead to Significantly More Warming



## Radiative Forcing, 1750-2010




Total net Forcing = + 2.3 w/m<sup>2</sup> (From 1.1 to 3.3 considering error bars)

Fig. 6.3 Radiative forcing caused by changes in the climate between 1750 and 2005. The error bars indicate the uncertainty of the estimate (based on values from Ciais et al., 2013).

# The Problem: AR5WG1\_CH. 9

- Cloud feedbacks represent the main cause for the range in modelled climate sensitivity (2.1C-4.7C).
  - the primary cloud-related factor contributing to the spread in equilibrium climate sensitivity is the low-level shortwave cloud feedback
  - Changes in the high-altitude clouds also induce climate feedbacks... the parameterisation of ice fall speed also emerges as one of the most important determinants of climate sensitivity

# Cloud Parameterizations

- Cloud processes occur in micron sized spaces. Cloud droplets are millimeters in diameter. (Hair is about 100 microns in diameter)
  - The climate model box size is currently ~100 km on a side.
  - You can not describe the physics occurring on micron sized objects if your smallest box is ~100 km on a side.
  - Therefore, you do not build your model cloud droplet by droplet from the fundamental physics. You use parameterizations which can differ from model to model.
- 

# Most Scenarios that Avoid the Most Dangerous Expected Effects of Climate Change Feature Overshoot in CO<sub>2</sub> and Later Drawdown.

- How to reduce climate impacts during this overshoot period?
- Some say Albedo Modification (see NAS Report)

*Paul J. Crutzen:*

## What to do?

- Artificially enhancing earth's albedo.
- Cool climate by adding sunlight reflecting aerosol in the stratosphere (Budyko, 1977; NAS, 1992) might again be explored...

# *Paul J. Crutzen*

## How to do it?

- Put SO<sub>2</sub> in the tropics at high altitude – It gets carried into stratosphere
- In the stratosphere, chemical and micro-physical processes convert SO<sub>2</sub> into sub-micrometer sulfate particles.
- This has been observed in volcanic eruptions e.g., Mount Pinatubo in June, 1991, which injected some 10 Tg S, initially as SO<sub>2</sub>, into the tropical stratosphere (Wilson et al., 1993; Bluth et al., 1992).

# Albedo Enhancement Schemes

- Add particles to stratosphere to increase albedo
- Enhance low-level, marine clouds to increase albedo
- Put a 1000 km x 1000 km parasol in space to reduce sunlight. (We do not know how to do this today.)
- First two are cheap compared to reducing emissions.

# Volcanic Eruptions Resulted in Long-lived Particles in the Stratosphere


- Temperature decrease following Pinatubo injection was 0.3C for 3 years with 20 Mt SO<sub>2</sub>.
- Robock and Liu [1994], models, and Trenberth and Dai [2007], observations following the 1991 Pinatubo eruption, found large reductions in the strength of the global hydrological cycle including in precipitation, soil moisture, and river flow.
- Possibly disrupts Asian Monsoon.
- Volcanic eruptions have also been observed to produce large stratospheric ozone depletion following the 1982 El Chichon and 1991 Pinatubo eruptions [Solomon, 1999].

- Not very expensive to inject  $\text{SO}_2$  gas. Atmospheric chemistry will make  $\text{H}_2\text{SO}_4$  which makes particles. Same chemistry as acid rain.
- If they are the same size as volcanic particles (around 1 micron) and reach the stratosphere, they will cool and persist.
- We do not know for sure that this will occur as hoped. The imagined injections do not mimic volcanic ones in location and amount.




- The injections imagined by the albedo enhancement schemes are occurring at 15-18 km altitude. The Pinatubo injection went in well above 20 km. The Albedo Enhancement Schemes assume that those different injection locations will result in the same particles in the end.

- I am skeptical. You note that we have measured volcanic, aircraft, natural particles at these altitudes and latitudes. My reasons for skepticism have to do with details of aerosol dynamics that are beyond the scope of this talk.



# Modeled Temperature Cools Quickly when Injections Start and Warms Quickly When They Stop (Robock et al., 2008). Models show decreased precipitation.



**Figure 1.** Global average surface air temperature change from the A1B anthropogenic forcing run (red), Arctic 3 Mt/a SO<sub>2</sub> (blue), tropical SO<sub>2</sub> 5 Mt/a (black), and tropical 10 Mt/a SO<sub>2</sub> (brown) cases in the context of the climate change of the past 125 years. Observations (green) are from the National Aeronautics and Space Administration Goddard Institute for Space Studies analysis [*Hansen et al., 1996*]

# THE BENEFITS, RISKS, AND COSTS OF STRATOSPHERIC GEOENGINEERING


A. Robock, A. Marquardt, B. Kravitz, and G. Stenchikov :GRL

## Benefits

1. Cool planet
2. Reduce or reverse sea ice melting
3. Reduce or reverse land ice sheet melting
4. Reduce or reverse sea level rise
5. Increase plant productivity
6. Increase terrestrial CO<sub>2</sub> sink

## Risks

1. Drought in Africa and Asia
2. Continued ocean acidification from CO<sub>2</sub>
3. Ozone depletion
4. No more blue skies
5. Less solar power
6. Environmental impact of implementation
7. Rapid warming if stopped




# THE BENEFITS, RISKS, AND COSTS OF STRATOSPHERIC GEOENGINEERING

A. Robock, A. Marquardt, B. Kravitz, and G. Stenchikov :GRL


## Risks, continued

8. Cannot stop effects quickly
9. Human error
10. Unexpected consequences
11. Commercial control
12. Military use of technology
13. Conflicts with current treaties
  14. Whose hand on the thermostat
  15. Ruin terrestrial optical astronomy
16. Moral hazard – the prospect of it working would reduce drive for mitigation
17. Moral authority – do we have the right to do this?

# Governance Questions

- Who gets to choose to do it?
  - Who gets to choose how much to do?
  - Who decides when to stop?
  - What kind of global agreements are necessary?
  - What if a company or country set up to try to do this?
- 


# Albedo enhancement of low-level maritime clouds (added)

- Put more droplets into marine stratus clouds by adding nuclei
    - More nuclei, smaller droplets, longer cloud persistence, whiter clouds and increased albedo
    - Some argue that we should be able to engineer the clouds and balance a doubling of CO<sub>2</sub>.
- 


Difficult to model these clouds and to predict outcomes. There are large uncertainties about impact. If brightening succeeds:

- Global changes in the distributions and magnitudes of ocean currents, temperature, rainfall and wind could result.
  - alter the land–ocean temperature contrast, since the radiative forcing produced would be only over the oceans.
- 

# Data Used to Apply 1<sup>st</sup> Law


- Satellite atmospheric temperatures (NOAA MSUs and NOAA AMSU- UAH and RSS analysis)
  - Ice sheet mass. GRACE
  - Ocean heat content: ARGO
  - Land mass temperatures: Bore Holes
- 

# Analysis

- First law of thermodynamics: Calculate changes in internal energy of each of these elements.
  - Masses, specific heats and heat of fusion are known.
  - Use measured changes in temperature and measured loss of land ice mass.
- 


# GRACE: Gravity Recovery and Climate Experiment

- 1.** If mass in a location changes (ice melts and water goes elsewhere) the gravity over that location will change.
- 2.** GRACE detects those changes in gravity and can therefore detect changes in mass distribution below.
- 3.** GRACE is beautiful. GRACE is physics.




- 5.** GRACE has been flying since March 2002 and has been used to study changes in ice, ground water and soil moisture
- 6.** GRACE is done. Follow-on GRACE is planned.

Velicogna, I., T. C. Sutterley, and M. R. van den Broeke (2014), Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data, *J. Geophys. Res. Space Physics*, 41, 8130–8137, doi:10.1002/2014GL061052.




Time series of ice mass,  $M(t)$ , in Gt (Gigatonne) for (a) the entire Greenland ice sheet (GIS), Blue: GRACE Total Mass Anomaly Red: Cumulative Surface Mass Balance (New snow - surface melting)

# Antarctic Ice Sheet – Mass Anomaly vs Time from GRACE





Time series of ice mass,  $M(t)$ , in Gt (Gigatonne) for (a) the entire Antarctic ice sheet (AIS), Blue: GRACE Mass Anomaly Red: Cumulative Surface Mass Balance (New snow - surface melting)

# ARGO Float





**Figure 1 | The typical cycle of an Argo float.** The float starts at the surface and dives to a depth of 1,000 m (the parking depth) where it rests for 9 to 10 days. After 9 days at rest it dives to a depth of 2,000 m, turns on

# ARGO Floats



# ARGO Data Alone






# Ocean heat Content From Argo and Earlier Systems

**Figure 5 | Argo estimates of global heat content.** Estimates of the heat content anomaly<sup>27</sup> for 60° S–20° S (black), 20° S–20° N (red) and 20° N–60° N (blue), and a global estimate (purple)<sup>42</sup>. Regression lines (dashed) are shown in the inset; adapted from ref. 27, Nature Publishing Group.

# Compare GAST and MSU Results



# DU Aerosol Group Develops Instruments, Makes In situ Aerosol Measurements and Does Analysis

NASA Global Hawk

Altitude Cruising Range: 45,000 – 65,000 ft

Duration: 33 hrs

DU Aerosol Payload

First Science Mission: Winter 2010



NASA WB-57F

Altitude Cruising Range: 45,000 – 65,000 ft


Duration: 6 hrs

DU Aerosol Payload in nose

# DU Aerosol Group: Instruments & Inlets, Missions & Science



NASA ER-2, DC-8, WB-57F  
NCAR C-130, HIAPER, Electra, UK FAAM, DLR Falcon  
Russian Geophysica, NPGS Twin Otter...



NASA Photo

